

**Table I: Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2006-2009**

County	Arthritis		Asthma		Heart Attack		Angina	
	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted
<b>OREGON</b>		<b>25.8%</b>		<b>9.7%</b>		<b>3.3%</b>		<b>3.4%</b>
<b>Baker</b>	31.6%	26.5%	7.9%	6.9%	3.5% †	2.7% †	6.2% †	4.9%
<b>Benton</b>	26.0%	27.9%	8.8%	9.4%	3.0%	3.3%	2.8%	3.0%
<b>Clackamas</b>	23.7%	22.4%	9.6%	9.7%	2.6%	2.5%	3.1%	2.9%
<b>Clatsop</b>	39.7%	39.1% *	7.9%	7.5%	4.8%	3.9%	4.0%	3.1%
<b>Columbia</b>	24.4%	22.4%	10.3%	9.8%	4.4% †	4.1% †	2.5% †	2.1% †
<b>Coos</b>	34.3%	28.4%	12.3%	13.1%	7.7% †	7.3% †	8.2%	7.7% †
<b>Crook</b>	25.2%	22.4%	5.7% †	5.6% †	—	2.0% †	—	—
<b>Curry</b>	31.5%	34.1% †	11.7%	12.6%	8.4%	4.4%	10.1%	6.0% *
<b>Deschutes</b>	29.9%	28.0%	9.0%	9.2%	3.0%	2.7%	2.8%	2.4%
<b>Douglas</b>	40.8%	36.8% *	12.6%	12.7%	5.7%	4.4%	5.2%	4.0%
<b>Grant</b>	18.3% †	11.2% †	12.0% †	13.6% †	4.2% †	2.9% †	3.6% †	2.5% †
<b>Harney</b>	30.6% †	26.8% †	12.9% †	13.3% †	7.5% †	6.0% †	—	—
<b>Hood River</b>	25.2% †	24.2%	2.8% †	—	6.2% †	5.8% †	3.9% †	3.6% †
<b>Jackson</b>	28.6%	25.4%	8.5%	8.3%	3.5%	2.9%	3.5%	2.9%
<b>Jefferson</b>	48.0%	47.5%	7.1% †	6.6% †	3.7% †	3.0% †	3.5% †	2.8% †
<b>Josephine</b>	31.2%	26.4%	11.3%	11.6%	4.9%	3.4%	5.7%	4.0%
<b>Klamath</b>	36.2%	32.5%	11.3%	11.5%	6.9%	6.0% *	4.7%	3.7%
<b>Lake</b>	29.8% †	25.2% †	6.3% †	7.2% †	—	—	—	—
<b>Lane</b>	28.3%	27.1%	10.5%	10.6%	3.4%	3.1%	3.8%	3.5%
<b>Lincoln</b>	31.8%	27.9%	10.7%	11.2%	5.6%	3.9%	3.4%	2.5% †
<b>Linn</b>	32.6%	29.5%	10.4%	10.5%	5.4%	4.5%	5.0%	4.2%
<b>Malheur</b>	27.3%	27.1%	8.4%	8.4%	3.7% †	3.4% †	2.0% †	1.7% †
<b>Marion</b>	25.1%	25.0%	11.4%	11.3%	4.1%	4.0%	3.9%	3.8%
<b>Morrow</b>	27.9% †	24.3% †	7.9% †	8.2% †	—	—	—	6.5% †
<b>Multnomah</b>	23.9%	24.3%	9.2%	9.2%	2.8%	2.9%	2.9%	3.0%
<b>Polk</b>	26.5%	25.6%	9.1%	9.1%	5.3%	5.2%	4.7%	4.4%
<b>Tillamook</b>	32.9%	28.7%	8.1% †	8.2% †	5.8% †	4.8% †	4.8% †	3.9% †
<b>Umatilla</b>	28.9%	27.8%	7.6%	7.5%	3.4%	3.0%	3.7%	3.4%
<b>Union</b>	31.9%	31.0%	12.8%	13.3%	4.3% †	4.0% †	6.2%	5.3%
<b>Wallowa</b>	45.8%	38.2%	8.0% †	6.2% †	—	—	8.7% †	5.4% †
<b>Washington</b>	22.2%	23.1%	9.0%	9.0%	2.4%	2.5%	2.9%	3.1%
<b>Yamhill</b>	24.6%	24.9%	11.5%	11.5%	3.2%	3.1%	3.2%	3.2%
<b>Gilliam/Wheeler</b>	—	—	—	—	—	6.3% †	—	—
<b>Sherman/Wasco</b>	20.7%	17.0%	6.1% †	6.8% †	6.0%	4.6%	6.0%	4.9% †

\* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

— This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2006-2009

**Table I (continued): Age-Adjusted and Unadjusted Prevalence of Selected Chronic Conditions among Adults, by County, Oregon 2006-2009**

County	Stroke		Diabetes		High Blood Pressure		High Blood Cholesterol	
	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted
<b>OREGON</b>		<b>2.3%</b>		<b>6.8%</b>		<b>25.8%</b>		<b>33.0%</b>
<b>Baker</b>	3.1% †	2.4% †	12.8%	11.4% *	39.0%	31.3%	56.2%	59.9% *
<b>Benton</b>	1.5%	1.6%	5.5%	6.3%	20.3%	22.9%	28.4%	25.8%
<b>Clackamas</b>	2.8%	2.6%	7.0%	6.6%	26.8%	25.6%	37.6%	33.1%
<b>Clatsop</b>	3.6%	2.8%	8.5%	7.1%	36.9%	33.9%	40.6%	28.3%
<b>Columbia</b>	3.1% †	2.8% †	7.3%	6.6%	34.0%	31.0%	41.6%	30.4%
<b>Coos</b>	5.4% †	5.7% †	12.6%	11.0%	34.1%	28.5%	45.5%	41.8%
<b>Crook</b>	—	—	9.9%	9.0%	48.0%	46.2% *	44.1%	41.8%
<b>Curry</b>	2.9% †	1.5% †	8.2%	6.1% †	40.3%	25.8%	46.9%	32.6%
<b>Deschutes</b>	1.4%	1.2% *	5.6%	5.0%	23.6%	20.6% *	36.0%	32.1%
<b>Douglas</b>	4.5%	3.8% *	11.4%	10.1% *	35.0%	31.4%	40.2%	35.1%
<b>Grant</b>	3.5% †	2.4% †	9.3% †	6.7%	33.4%	21.7%	49.2%	42.1% †
<b>Harney</b>	—	—	9.5% †	8.1% †	28.8%	22.2%	48.1%	37.3%
<b>Hood River</b>	—	—	4.9% †	4.4% †	25.5%	23.2%	17.7% †	13.4% †
<b>Jackson</b>	2.6%	2.2%	7.2%	6.4%	29.2%	26.8%	41.5%	35.5%
<b>Jefferson</b>	2.3% †	1.9% †	7.3%	6.5%	20.5%	16.9% *	29.4%	20.2% †
<b>Josephine</b>	3.9%	3.3%	8.2%	5.8%	38.0%	34.3%	45.5%	43.7%
<b>Klamath</b>	2.5% †	2.3% †	7.9%	7.3%	32.0%	29.4%	38.9%	34.3%
<b>Lake</b>	—	—	—	—	39.6%	30.2%	58.6%	45.1%
<b>Lane</b>	2.4%	2.2%	6.5%	6.3%	28.7%	27.3%	39.2%	33.6%
<b>Lincoln</b>	3.4% †	2.8% †	11.3%	8.9%	39.8%	36.2%	54.7%	41.0%
<b>Linn</b>	4.0%	3.7%	9.0%	7.9%	29.7%	27.7%	34.4%	29.6%
<b>Malheur</b>	2.7% †	2.6% †	8.0%	7.6%	30.2%	27.8%	41.2%	37.5%
<b>Marion</b>	2.4%	2.3%	8.3%	8.3%	28.1%	27.9%	32.1%	29.1%
<b>Morrow</b>	—	—	7.1% †	6.8% †	19.9% †	16.1%	33.8% †	23.1% †
<b>Multnomah</b>	1.8%	1.8%	6.0%	6.2%	22.8%	23.0%	36.4%	33.7%
<b>Polk</b>	2.5%	2.2%	6.6%	6.0%	25.9%	24.4%	34.7%	31.0%
<b>Tillamook</b>	2.8% †	2.0% †	10.0%	8.3%	35.9%	32.2%	40.9%	29.6%
<b>Umatilla</b>	2.9%	2.7%	9.6%	9.3%	32.1%	30.9%	46.1%	39.7%
<b>Union</b>	4.4% †	3.9% †	6.5%	6.5%	25.5%	22.6%	41.6%	36.1%
<b>Wallowa</b>	—	—	4.0% †	2.4% †	30.3%	18.8%	44.4%	32.4%
<b>Washington</b>	1.8%	1.9%	5.6%	5.9%	21.8%	22.9%	32.5%	30.2%
<b>Yamhill</b>	2.2%	2.2%	7.5%	7.5%	27.2%	26.8%	36.1%	34.9%
<b>Gilliam/Wheeler</b>	—	—	15.6% †	12.4% †	—	—	—	—
<b>Sherman/Wasco</b>	2.7% †	1.9% †	6.0%	5.3%	31.4%	27.8%	34.0%	26.8%

\* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

— This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are usually adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+), however, the Grant, Jefferson and Morrow high blood cholesterol estimates were based on collapsed age groups (18-54 and 55+) due to small numbers.

Source: Oregon BRFSS County Combined Dataset 2006-2009