

**Table III: Age-Adjusted and Unadjusted Prevalence of Preventive Health Screening among Adults, by County, Oregon 2006-2009**

County	% who had their cholesterol checked within past five years (≥ 18 years old)		% who had a PAP test within past three years (women 18-65 years old with intact cervix)	
	unadjusted	age-adjusted	unadjusted	age-adjusted
<b>OREGON</b>		<b>71.3%</b>		<b>85.8%</b>
Baker	72.8%	66.8%	76.0%	75.8%
Benton	70.3%	75.1%	97.0%	95.1% *
Clackamas	72.3%	70.6%	88.7%	88.3%
Clatsop	72.8%	68.4%	56.4%	61.3% *
Columbia	72.3%	67.9%	89.6%	89.8%
Coos	80.9%	75.1%	78.5%	78.3%
Crook	83.0%	81.7%	—	—
Curry	65.6%	49.7% *	—	—
Deschutes	76.2%	73.3%	80.3%	79.8%
Douglas	77.8%	73.6%	80.1%	79.9%
Grant	76.7%	54.1% *	—	—
Harney	67.4%	58.6%	—	—
Hood River	69.9%	69.0%	94.8%	92.8%
Jackson	73.9%	71.1%	82.2%	85.5%
Jefferson	58.9%	56.3% *	64.8%	63.1% *
Josephine	79.7%	75.3%	81.7%	81.6%
Klamath	74.4%	72.2%	81.2%	81.1%
Lake	74.5%	69.6%	—	—
Lane	68.5%	67.3%	85.2%	86.1%
Lincoln	65.6%	54.1% *	82.9%	85.1%
Linn	70.6%	68.1%	83.3%	83.3%
Malheur	76.4%	74.8%	89.8%	88.4%
Marion	73.1%	73.0%	82.7%	84.3%
Morrow	53.0%	44.5% *	—	—
Multnomah	72.0%	71.6%	86.6%	86.6%
Polk	73.5%	72.1%	77.1%	77.6%
Tillamook	76.1%	70.1%	86.8%	87.9%
Umatilla	73.6%	73.2%	82.6%	80.3%
Union	71.9%	70.2%	89.8%	89.9%
Wallowa	82.2%	65.2% *	—	—
Washington	73.8%	74.0%	91.5%	91.5%
Yamhill	70.7%	71.4%	83.9%	83.2%
Gilliam/Wheeler	—	—	—	—
Sherman/Wasco	68.0%	63.1%	—	—

\* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

— This number is suppressed because it is statistically unreliable.

Age-adjusted estimates are adjusted to the 2000 Standard Population using three age groups (18-34, 35-54, and 55+).

Source: Oregon BRFSS County Combined Dataset 2006-2009

**Table III (continued): Age-Adjusted and Unadjusted Prevalence of Preventive Health Screening among Adults, by County, Oregon 2006-2009**

County	% who had a mammogram within past two years (women 50-74 years old)	% who had an FOBT <sup>^</sup> in the past year or a Colonoscopy/ Sigmoidoscopy within the past five years (50-75 year olds)
<b>OREGON</b>	<b>81.9%</b>	<b>56.8%</b>
Baker	70.6%	47.0%
Benton	89.5%	67.9%
Clackamas	86.4%	57.2%
Clatsop	76.3%	44.1%
Columbia	84.3%	62.2%
Coos	89.6%	62.2%
Crook	—	47.3%
Curry	65.9%	61.8%
Deschutes	72.8%	58.9%
Douglas	77.3%	60.2%
Grant	—	58.5%
Harney	—	35.4%†
Hood River	—	26.3%†
Jackson	82.4%	54.5%
Jefferson	—	56.1%
Josephine	72.2%	48.5%
Klamath	75.2%	47.4%
Lake	—	—
Lane	82.8%	60.6%
Lincoln	63.8%*	46.7%
Linn	77.4%	57.8%
Malheur	70.5%	42.6%
Marion	85.9%	55.4%
Morrow	—	—
Multnomah	85.7%	57.8%
Polk	89.0%	59.9%
Tillamook	72.2%	46.0%
Umatilla	81.0%	53.6%
Union	71.8%	58.4%
Wallowa	—	56.1%
Washington	86.5%	58.2%
Yamhill	76.3%	59.6%
Gilliam/Wheeler	—	—
Sherman/Wasco	80.0%	56.8%

\* Statistically significant difference compared with Oregon.

† This number may be statistically unreliable and should be interpreted with caution.

— This number is suppressed because it is statistically unreliable.

<sup>^</sup> Fecal Occult Blood Test

Source: Oregon BRFSS County Combined Dataset 2006-2009